



GeneInact seq (2000)-2.ST25.txt
SEQUENCE LISTING

<110> GMR Epigenetics Corporation
<120> Gene inactivation by targeted DNA methylation
<130> GMR-001ORD
<140> US 09/643,128
<141> 2000-08-21
<150> US 60/196,749
<151> 2000-04-12
<150> US 60/214,148
<151> 2000-06-26
<160> 54
<170> PatentIn version 3.3
<210> 1
<211> 11
<212> DNA
<213> Artificial
<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(11)
<223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions 7 and 10.

2). Semi-methylated hairpin loop

<220>
<221> CpG dinucleotide
<222> (1)..(11)

<400> 1
cgacgtcgtc g

11

<210> 2
<211> 22
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(22)
<223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions 5, 9 and 20

2). Semi-methylated hairpin loop

<220>

<221> CpG dinucleotide

<222> (1)..(22)

<400> 2

agcccgggct gggaggagtc gg

22

<210> 3

<211> 33

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(33)

<223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #7, #10, #16, #20, #31.

<220>

<221> CpG dinucleotide

<222> (1)..(33)

<223> 2) Semi-methylated loop

<220>

<221> CpG dinucleotide

<222> (1)..(33)

<400> 3

cgacgtcgtc gagcccgggc tgggaggagt cgg

33

<210> 4

<211> 21

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(21)

<223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #12 and #19.

2). Sequence derived from the promoter of human c-myc gene.

3). Semi-methylated loop.

<220>

<221> CpG dinucleotide

<222> (1)..(21)

GeneInact seq (2000)-2.ST25.txt

<220>

<221> CpG dinucleotide

<222> (1)..(21)

<400> 4

tcgctaattct ccgcccaccg g

21

<210> 5

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine at positions 3 and 18.

2) Complementary to promoter of human c-myc.

3) Semi-methylated hairpin.

<400> 5

accggccctt tataatgcga

20

<210> 6

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #3 and #10.

2). Sequence derived from the promoter of human c-myc gene.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions 3 and 10. 3

2) Sequence complementary to the promoter of human c-myc gene.

3) Semi-methylated hairpin.

<400> 6
tccgcccacc ggccctttat

20

<210> 7
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #3, #10 and #16.
2). Sequence derived from the promoter of HIV gene.

<220>
<221> CpG dinucleotide
<222> (1)..(18)

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 3). Semi-methylated hairpin.

<400> 7
cacgtagccc gagagctg

18

<210> 8
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #7, #9 and #16.
2). Sequence derived from the promoter of HIV gene.

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 3). Semi-methylated hairpin.

<220>
<221> CpG dinucleotide
<222> (1)..(18)

<400> 8
 cccgagagct gcatccgg

18

<210> 9
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1): Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #13 and #16.
 2). Sequence derived from the promoter of HIV gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<400> 9
 gctgcatata agcagctg

18

<210> 10
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #4, #10 and #15.
 2) Sequence derived from the promoter of human uPAR gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin.

<400> 10

aggcgccac gcatctgg

<210> 11
 <211> 19
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #10 and #17.
 2). Sequence derived from the promoter of human uPAR gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 3). Semi-methylated hairpin.

<400> 11
 tcgctctttc gcaaaacgt

<210> 12
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions 2, 7 and 13.
 2). Sequence derived from the promoter of human uPAR gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin.

<400> 12
 acgcatctgg ggctgact

GeneInact seq (2000)-2.ST25.txt

<210> 13
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #10 and #16.
2). Sequence derived from the promoter of human VEGF gene.

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 3). Semi-methylated hairpin loop.

<400> 13
gttataaatc gccccgc

18

<210> 14
<211> 23
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(23)
<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2 and #21.
2). Sequence derived from the promoter of human VEGFR gene.

<220>
<221> CpG dinucleotide
<222> (1)..(23)
<223> 3). Semi-methylated hairpin loop

<400> 14
gctggggaaa gggtataaat cgc

23

<210> 15
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human

sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #10 and #16.
 2). Sequence derived from the promoter of human VEGFR gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin loop

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<400> 15
 accccttgac gtcaccag

18

<210> 16
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #7, #14 and #16.
 2). Sequence derived from the promoter of human VEGFR gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin loop

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<400> 16
 cttcatcgag gtccgcgg

18

<210> 17
 <211> 17
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(17)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #7 and #15.
 2). Sequence derived from the promoter of human flk-1 gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(17)

<220>
 <221> CpG dinucleotide
 <222> (1)..(17)
 <223> 3). Semi-methylated hairpin loop

<400> 17
 cctgcactga gtcccgg

17

<210> 18
 <211> 21
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(21)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2 and #19.
 2). Sequence derived from the promoter of human VEGF gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(21)

<220>
 <221> CpG dinucleotide
 <222> (1)..(21)
 <223> 3). Semi-methylated hairpin loop

<400> 18
 acgggagagc ccctcctccg c

21

<210> 19
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

GeneInact seq (2000)-2.ST25.txt

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #11 and #15.

2). Sequence derived from the promoter of human integrin beta 3 gene.

<220>
<221> CpG dinucleotide
<222> (1)..(18)

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 3). Semi-methylated hairpin loop

<400> 19
cactgtgggg cgggcgga

18

<210> 20
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #3, #15 and #17.

2). Sequence derived from the promoter of human integrin beta 3 gene.

<220>
<221> CpG dinucleotide
<222> (1)..(18)

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 3). Semi-methylated hairpin loop

<400> 20
tgcgtcccac ccaccgcg

18

<210> 21
<211> 19
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2 and #9.
 2). Sequence derived from the promoter of human 12-lipoxygenase gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 3). Semi-methylated hairpin loop

<400> 21
 ccgcagaccg gtcctttaa

19

<210> 22
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #7 and #13.
 2). Sequence derived from the promoter of human 12-lipoxygenase gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin loop

<400> 22
 cctgggcggt cccgggca

18

<210> 23
 <211> 19
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human

sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at sequence positions #4, #7 and #16.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 2). Sequence derived from the promoter of human beta-amyloid protein precursor gene.

3). Semi-methylated hairpin loop.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)

<400> 23
 ctccgtcagt ttctcggc

19

<210> 24
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) positions #3, #6, #12 and #15.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 2). Sequence derived from the promoter of human 12-lipoxygenase gene.

3). Semi-methylated hairpin loop.

<400> 24
 atcagctgac tcgcctgg

18

<210> 25
 <211> 19
 <212> DNA
 <213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #4, #14 and #16.

2). Sequence derived from the promoter of human VEGF gene.

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<223> 3). Semi-methylated hairpin loop.

<400> 25

tagcggggag gatcgcgga

19

<210> 26

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #8, #11 and #18.

2). Sequence derived from the promoter of human VEGF gene.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 3). Semi-methylated hairpin loop.

<400> 26

taaaagtcgg ctggtagcgg

20

<210> 27

<211> 18

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human

sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #2.
 2). Sequence derived from the promoter of human IGF1 gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin loop.

<400> 27
 tctgtgctct agttttaa

18

<210> 28
 <211> 18
 <212> DNA
 <213> Artificial

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #5 and #13.
 2). Sequence derived from the promoter of human IGF1 gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin loop.

<400> 28
 ccagctgttt tcctgtct

18

<210> 29
 <211> 20
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

GeneInact seq (2000)-2.ST25.txt

<220>
 <221> CpG dinucleotide
 <222> (1)..(20)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #2.
 2). Sequence derived from the promoter of human Her2 gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(20)

<220>
 <221> CpG dinucleotide
 <222> (1)..(20)
 <223> 3). Semi-methylated hairpin loop.

<400> 29
 gctgcttgag gaagtataag 20

<210> 30
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(20)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #18.
 2). Sequence derived from the promoter of human Her2 gene.
 3). Semi-methylated hairpin loop.

<400> 30
 agaataagtg tgtgaagctg 20

<210> 31
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #4.

2). Sequence derived from the promoter of human TNF-alpha gene.

<220>

<221> CpG dinucleotide

<222> (1)..(18)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #4.

2). Sequence derived from the promoter of human TNF-alpha gene.

3). Semi-methylated hairpin loop.

<400> 31

tgccgttcct ctataaag

18

<210> 32

<211> 18

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(18)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #7, #12 and #16.

2). Sequence derived from the promoter of human TNF-alpha gene.

<220>

<221> CpG dinucleotide

<222> (1)..(18)

<220>

<221> CpG dinucleotide

<222> (1)..(18)

<223> 3). Semi-methylated hairpin loop.

<400> 32

agggacctga gcgtccgg

18

<210> 33

<211> 21

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(21)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2 and #7.

2). Sequence derived from the promoter of human TNF-beta gene.

3). Semi-methylated hairpin loop.

<220>

<221> CpG dinucleotide

<222> (1)..(21)

<400> 33

tcgccccagg gacatataaa g

21

<210> 34

<211> 18

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(18)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #11.

2). Sequence derived from the promoter of human TNF-beta gene.

3). Semi-methylated hairpin loop.

<400> 34

catataaagg cagttggt

18

<210> 35

<211> 18

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(18)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #4, #11 and #15.

2). Sequence derived from the promoter of human TNF-beta gene.

<220>

<221> CpG dinucleotide

<222> (1)..(18)

<220>

<221> CpG dinucleotide

<222> (1)..(18)

<223> 3). Semi-methylated hairpin loop.

<400> 35
 acccagccag cagacgct

18

<210> 36
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2 and #8.

2). Sequence derived from the promoter of human interleukin 4 gene.

3). Semi-methylated hairpin loop.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)

<400> 36
 tcggtttcag caatttga

18

<210> 37
 <211> 19
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #16.

2). Sequence derived from the promoter of human IL-4 gene.

3). Semi-methylated hairpin loop.

<400> 37
 tagagatatc tttgtcagc

19

<210> 38
 <211> 19
 <212> DNA
 <213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #3.

2). Sequence derived from the promoter of human GM-CSF gene.

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<223> 3). Semi-methylated hairpin loop.

<400> 38

ctctgtgtat ttaagagct

19

<210> 39

<211> 19

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2 and #9.

2). Sequence derived from the promoter of human GM-CSF gene.

3). Semi-methylated hairpin loop.

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<400> 39

ccgcctccct ggcattttg

19

<210> 40

<211> 19

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

GeneInact seq (2000)-2.ST25.txt

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #2.
 2). Sequence derived from the promoter of human IL-2 gene.
 3). Semi-methylated hairpin loop.

<400> 40
 ccagagagaa gagtataat 19

<210> 41
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #5.
 2). Sequence derived from the promoter of human bcl-2 gene.
 3). Semi-methylated hairpin loop.

<400> 41
 atagctggat tataactc 18

<210> 42
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #2.
 2). Sequence derived from the promoter of human bcl-2 gene.
 3). Semi-methylated hairpin loop.

<400> 42
 tcgtccaaga atgcaaag 18

<210> 43

GeneInact seq (2000)-2.ST25.txt

<211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #13.
 2). Sequence derived from the promoter of human HBV gene.
 3). Semi-methylated hairpin loop.

<400> 43
 cagccatgga aacgatgt

18

<210> 44
 <211> 19
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #6 and #17.
 2). Sequence derived from the promoter of human HBV gene.
 3). Semi-methylated hairpin loop.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)

<400> 44
 tgaagcgaag tgcacacgg

19

<210> 45
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide

GeneInact seq (2000)-2.ST25.txt

<222> (1)..(18)
<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #4, #13 and #15.

2). Sequence derived from the promoter of human HBV gene.

<220>
<221> CpG dinucleotide
<222> (1)..(18)
<223> 3). Semi-methylated hairpin loop.

<400> 45
agacggtgag accgcgta

18

<210> 46
<211> 19
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(19)
<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at position #16.

2). Sequence derived from the promoter of human HBV gene.

3). Semi-methylated hairpin loop.

<400> 46
tgcattggtgc tgggtgcgca

19

<210> 47
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> This sequence is complementary to and methylates a human sequence.

<220>
<221> CpG dinucleotide
<222> (1)..(20)
<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #5, #12 and #18.

2). Sequence derived from the promoter of CMV gene.

<220>
<221> CpG dinucleotide
<222> (1)..(20)
<223> 3). Semi-methylated hairpin loop.

<400> 47

tgggcggtag gcgtgtacgg

<210> 48
 <211> 19
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #14 and #17.
 2). Sequence derived from the promoter of CMV gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 3). Semi-methylated hairpin loop.

<400> 48
 acggtaaagt gcccgcctg

<210> 49
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2 and #13.
 2). Sequence derived from the promoter of CMV gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin loop.

<400> 49
 gcgtcaatgg ggcggagt

<210> 50
 <211> 20
 <212> DNA
 <213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2 and #17.

2). Sequence derived from the promoter of human c-fos gene.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 3). Semi-methylated hairpin loop.

<400> 50

acgcttgta taaaagcagt

20

<210> 51

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #13 and #18.

2). Sequence derived from the promoter of human c-fos gene.

<220>

<221> CpG dinucleotide

<222> (1)..(20)

<223> 3). Semi-methylated hairpin loop.

<400> 51

tcgtactcca accgcatctg

20

<210> 52

<211> 19

<212> DNA

<213> Artificial sequence

<220>

<223> This sequence is complementary to and methylates a human sequence.

<220>

<221> CpG dinucleotide

<222> (1)..(19)

<223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #15 and #17.

2). Sequence derived from the promoter of human raf-1 gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(19)
 <223> 3). Semi-methylated hairpin loop.

<400> 52
 ccgagagtct taatcgcg

19

<210> 53
 <211> 18
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 1). Phosphothioate oligonucleotide modified with 5-methylcytidine (m5C) at positions #2, #4 and #12.

2). Sequence derived from the promoter of human raf-1 gene.

<220>
 <221> CpG dinucleotide
 <222> (1)..(18)
 <223> 3). Semi-methylated hairpin loop.

<400> 53
 tcgcgagaa tcggaggc

18

<210> 54
 <211> 22
 <212> DNA
 <213> Artificial sequence

<220>
 <223> This sequence is complementary to and methylates a human sequence.

<300>
 <301> Yao X, Hu JF, Daniels M, Shiran H, Zhou X, Yan H, Lu H, Zeng Z, Wang Q, Li T, Hoffman AR.
 <302> A methylated oligonucleotide inhibits IGF2 expression and enhances survival in a model of hepatocellular carcinoma.
 <303> J Clin Invest
 <304> 111
 <305> 2
 <306> 265-273
 <307> 2003-01-15
 <308> 12531883
 <309> 2003-03-15
 <313> (1)..(22)

<400> 54

ggtcacggtc aggcgtagat gg GeneInact seq (2000)-2.ST25.txt

22